

On 3d extensions of AGT relation

D.Galakhov*, A.Mironov†, A.Morozov‡, A.Smirnov§

ABSTRACT

An extension of the AGT relation from two to three dimensions begins from connecting the theory on domain wall between some two S-dual SYM models with the 3d Chern-Simons theory. The simplest kind of such a relation would presumably connect traces of the modular kernels in 2d conformal theory with knot invariants. Indeed, the both quantities are very similar, especially if represented as integrals of the products of quantum dilogarithm functions. However, there are also various differences, especially in the "conservation laws" for integration variables, which hold for the monodromy traces, but not for the knot invariants. We also discuss another possibility: interpretation of knot invariants as solutions to the Baxter equations for the relativistic Toda system. This implies another AGT like relation: between 3d Chern-Simons theory and the Nekrasov-Shatashvili limit of the 5d SYM.

1 Introduction

In [1] (see also [2]-[12]) a long expected relation was suggested, which can be considered as one of possible 3+3 counterparts of the celebrated 2+4 AGT relation [13] between conformal blocks and Nekrasov functions. This new relation is supposed to identify the modular transformation kernels $M(a, a')$ of conformal blocks and matrix elements in 3d Chern-Simons theory. In its simplest version, relation is between the trace of the modular kernel, considered as a function of external dimensions, and Hikami integrals [14], representing the Chern-Simons partition functions on S^3/K (a 3d sphere with a knot K removed) considered as functions of monodromies around K . A modest task of this letter is to discuss possible formulations of such a relation in an explicit form, leaving aside all the general context and reasoning, discussed in great detail in the profound text [1]. We point out some problems with exact identification of the modular trace and the knot invariants. We also attract attention to another similarity: between the knot invariants and 5d Seiberg-Witten theory, which implies still another kind of AGT relation, involving the 3d Chern-Simons theory.

2 Modular kernel

The conformal block $B_\Gamma(a|m|q)$ for a given graph Γ depends on three kinds of variables: a and m are parameters (α -parameters) on internal lines and external legs respectively (the corresponding conformal dimensions are quadratic in these parameters), and q parameterizes the graph itself. The modular transformation does not change the graph Γ , while changing $q \rightarrow q'$ and permuting the entries within the set α of external-legs parameters and is realized as an integral transform in a variables:

$$B_\Gamma(a|\alpha|q) = \int M(a, a') B_\Gamma(a'|\alpha'|q') d\mu(a') \quad (1)$$

The function $M(a, a')$ depends on Γ and α , but not on q , and is called the modular kernel, associated with the particular modular transformation $q \rightarrow q'$. In spirit, it is a Fourier kernel:

$$M(a, a') \sim \exp\left(\frac{4\pi i a a'}{\epsilon_1 \epsilon_2} + \dots\right) \quad (2)$$

where the corrections are less singular at small ϵ_1 and ϵ_2 .

*ITEP, Moscow, Russia and MIPT, Dolgoprudny, Russia; galakhov@itep.ru

†Lebedev Physics Institute and ITEP, Moscow, Russia; mironov@itep.ru; mironov@lpi.ru

‡ITEP, Moscow, Russia; morozov@itep.ru

§ITEP Moscow, Russia and MIPT, Dolgoprudny, Russia; asmirnov@itep.ru




Figure 1: Toric conformal block diagram, $\Delta_a = \frac{\epsilon^2/4+a^2}{\epsilon_1\epsilon_2}$, $\Delta_\alpha = \frac{\epsilon^2/4+\alpha^2}{\epsilon_1\epsilon_2}$.

The simplest example is provided by the modular transformation of the toric 1-point function, Fig.1 which on the gauge theory side describes the theory with adjoint matter with mass $m = -i\alpha + \frac{\epsilon}{2} = -2i\tilde{\alpha}$:

$$\mathcal{B}(a|\alpha|\tau) = \int \mathcal{M}(a, a'|\alpha) \mathcal{B}(a'|\alpha|\tau) d\mu(a') \quad (3)$$

Here [15]

$$\mathcal{B}(a|\alpha|\tau) = q^{-(\nu+1)/24} \eta(q)^\nu e^{\frac{2\pi i \tau a^2}{\epsilon_1 \epsilon_2}} \left(1 + 2q \frac{(\epsilon_1 - m)(\epsilon_2 - m)}{\epsilon_1 \epsilon_2} \frac{(\epsilon^2 - 4a^2 + m(m - \epsilon))}{(\epsilon^2 - 4a^2)} + O(q^2) \right) \quad (4)$$

with $\nu = 1 - \frac{2m(\epsilon-m)}{\epsilon_1 \epsilon_2}$, $\epsilon = \epsilon_1 + \epsilon_2$ and $\eta(q) = q^{\frac{1}{24}} \prod_{k=1}^{\infty} (1 - q^k)$, $q = e^{2\pi i \tau}$. According to [16] the modular kernel is given by

$$\mathcal{M}(a, a'|\alpha) = \frac{2^{3/2}}{s(\alpha)} \int \frac{s(a+r+\tilde{\alpha})s(a-r+\tilde{\alpha})}{s(a+r-\tilde{\alpha})s(a-r-\tilde{\alpha})} e^{\frac{4\pi i r a'}{\epsilon_1 \epsilon_2}} dr \quad (5)$$

where the measure is

$$d\mu(a') = 4 \sinh(2\pi \epsilon_1 a') \sinh(2\pi \epsilon_2 a') da' \quad (6)$$

The function $s(z)$ is the "quantum dilogarithm" [17, 18], the ratio of two digamma-functions [17],

$$s(z|\epsilon_1, \epsilon_2) \sim \prod_{m,n \geq 0} \frac{(m + \frac{1}{2}) \epsilon_1 + (n + \frac{1}{2}) \epsilon_2 - iz}{(m + \frac{1}{2}) \epsilon_1 + (n + \frac{1}{2}) \epsilon_2 + iz} \quad (7)$$

Note that $\mathcal{M}(a, a'|\alpha)$ depends on the external momentum $\tilde{\alpha}$, the internal dimension $\Delta(\alpha)$ and the central charge $c = 1 + 6 \frac{\epsilon^2}{\epsilon_1 \epsilon_2}$, but not on the modular parameter τ .

When $\alpha \rightarrow 0$, the conformal block becomes pure classical: $\mathcal{B}(a|\alpha|\tau) \rightarrow \frac{1}{\eta(q)} e^{2i\pi \tau a^2 / \epsilon_1 \epsilon_2}$ (the last factor in the brackets in (4) turns into $q^{1/12} \eta^{-2}(q)$ when $m = 0$), and the modular transformation turns into the ordinary Fourier transform:

$$\mathcal{M}(a, a'|0) \mu'(a) \xrightarrow{\tilde{\alpha} = i\epsilon/2} \sqrt{2} \cos \left(4\pi i \frac{aa'}{\epsilon_1 \epsilon_2} \right) \quad (8)$$

so that indeed

$$\frac{e^{-2\pi i \tau^{-1} a^2 / (\epsilon_1 \epsilon_2)}}{\eta(-\tau^{-1})} = \int \frac{da'}{\sqrt{\epsilon_1 \epsilon_2}} e^{4\pi i a a' / (\epsilon_1 \epsilon_2)} \frac{e^{2\pi i \tau a'^2 / (\epsilon_1 \epsilon_2)}}{\eta(\tau)} \quad (9)$$

It is a somewhat non-trivial exercise to deduce (8) from eq.(5), see Appendix A for the simplified case $\epsilon_1 = -\epsilon_2 = g_s$.

Notice that the monodromy kernel satisfies the unitarity relation

$$\int d\mu(a) M(a, b) M^*(a, b') = \frac{d\mu(b')}{db'} \delta(b - b') \quad (10)$$

Therefore, it is natural to define the trace as $\text{Tr} \sim \int \frac{d\mu(a)}{d\mu(a')} da' \delta(a - a')$. Thus defined trace of the monodromy kernel (5) contains two integrals, but they actually split:

$$\int \mathcal{M}(a, a|\alpha) da = \frac{2^{3/2}}{s(\alpha)} \int \int \frac{s(a+r+\tilde{\alpha})s(a-r+\tilde{\alpha})}{s(a+r-\tilde{\alpha})s(a-r-\tilde{\alpha})} e^{4\pi i r a} dr da = \frac{2^{3/2}}{s(\alpha)} T_+(\tilde{\alpha}) T_-(\tilde{\alpha}) \quad (11)$$

where

$$T_{\pm}(\tilde{\alpha}) = \int \frac{s(z+\tilde{\alpha})}{s(z-\tilde{\alpha})} e^{\pm i\pi z^2} dz \quad (12)$$

In fact, the quantities like $T_{\pm}(\tilde{\alpha})$ are well known from Chern-Simons theory, and this opens a way towards $3d$ extensions of the AGT conjecture.

3 Exempts from the knot theory

The polynomial knot invariants can be defined as averages of the Wilson loop along the knot in the topological Chern-Simons (CS) theory [19]:

$$\langle K \rangle_R = \left\langle \text{Tr}_R P \exp \oint_K A \right\rangle_{CS} \quad (13)$$

This invariant depends on the knot K , Lie algebra G , its representation R , the CS coupling constant $\hbar = \log q = \frac{2\pi i}{k}$ (sometimes k is shifted to $k + C_A$, as in the WZNW model [20]), and, additionally, on the monodromy u , which describes deviation from the periodicity of the field A while circling around the knot. One may think of u as of the eigenvalue of $PSL(2)$ monodromy matrix around the knot K . On the other hand, one may think that u takes value in the Cartan subalgebra of the gauge group $SU(2)$, and, in this way, describes the representation R living on the knot K .

The averages $\langle K \rangle_R$ are clever generalizations of the ordinary characters and, as all exact correlators, possess hidden integrability properties [21, 25]. As a manifestation of this hidden structure, the averages $\langle K \rangle_R$ satisfy K -dependent difference equations in the $R(!)$ variable [26], which allows one to consider them as belonging to the family of generalized q -hypergeometric series. A $q \rightarrow 1$ limit of these equations defines the spectral curve $\Sigma(K)$ and the saddle point of the corresponding integral representation defines the associated Seiberg-Witten (SW) differential. After that, the full $\hbar$ -dependence can be reconstructed with the help of the AMM/EO topological recursion [27] from this SW data [9].

A nice property of invariants (13) is that at $u = 0$ they are finite polynomials in $q = \exp(\frac{2\pi i}{k})$. In the literature, these polynomials normalized by the quantum dimension have different conventional names corresponding to the choice of the group and representation. Here, for convenience we represent a table explaining the correspondence:

Group\Representation	Fundamental representation	General representation with weight λ
$N = 0$	Convey-Alexander polynomial	—
$SU(2)$	Jones polynomial	Colored Jones polynomial
$SU(N)$	HOMFLY polynomial	Colored HOMFLY polynomial
$SO(N)$	Kauffman polynomial	Colored Kauffman polynomial
$\{SU(N)\}_t$	superpolynomial	Colored superpolynomial

The last line of this table describes an extension from (quantum) groups to MacDonald characters which leads to a one-parametric deformation (t -deformation) of (13), to superpolynomials [22, 23] involving the Khovanov homology [24]. Further extension from MacDonald to the Askey-Wilson-Kerov level remains untouched so far.

Six kinds of representations are currently known for the Wilson averages in Chern-Simons theory, we briefly describe them in the following subsections and provide some explicit examples in Appendix B.

3.1 Representation through quantum R -matrix [28]

This representation appears when calculations in CS theory are done in the temporal gauge $A_0 = 0$ [29]. Then the propagator is ultralocal, and only the crossings (c) and extremal (e) points in projection of K on the xy plane contribute, and the answer can be schematically written as

$$\langle K \rangle_R = \text{Tr}_R \overrightarrow{\prod_{e,c}} \mathcal{U}_e \mathcal{R}_c \quad (14)$$

where the ordered product is taken along the line K , $\mathcal{R}$ is the quantum R -matrix in representation R , and $\mathcal{U} = q^\rho$ is the "enhancement" of the $\mathcal{R}$ -matrix in the same representation. For the braid representation of the knot, this formula reduces to the well-known formula for the quantum group invariants of knot:

$$\langle K \rangle_R = q^{-w(b_K) \Omega_2(R)} \text{qtr}_R(b_K) \quad (15)$$

Here $b_K \in B_n$ is the element of the braid group, representing the knot K , i.e. its closure gives K , $\Omega_2(R) = \text{tr}_R(T^a T^a) / \dim R$ is the value of quadratic Casimir function in the representation R and qtr_R is the quantum trace over the representation R , $\text{qtr}_R b_K = \text{tr}_R b^K q^{\rho^{\otimes n}}$. The function $w(b_K)$ is the so called writhe number of the braid b_K , it is equal to the total sum of orientations of the crossings:

$$b_K = \prod_{\{k\}} g_k^{n_k} \implies w(b_K) = \sum_{\{k\}} n_k$$

Calculation of (15) for some particular knot is just a matter of multiplication and taking a trace of relatively big matrices. The braid representations of several first knots and the writhe numbers of the corresponding closures are summarized in the following table:

Knot	Braid representation	Writhe
3 ₁	$b = g_1^3 \in B_2$	$w(\hat{b}) = 3$
4 ₁	$b = g_2 g_1^{-1} g_2 g_1^{-1} \in B_3$	$w(\hat{b}) = 1$
5 ₁	$b = g_1^5 \in B_2$	$w(\hat{b}) = 5$
5 ₂	$b = g_2^3 g_1 g_2^{-1} g_1 \in B_3$	$w(\hat{b}) = 4$
6 ₁	$b = g_1 g_2^{-1} g_3 g_1 g_2^{-1} g_3^{-2} \in B_4$	$w(\hat{b}) = -1$

(16)

3.2 Representation through classical R -matrix and quantum associator [30]

A similar representation for $\langle K \rangle_R$ through classical instead of quantum R -matrices appears in the calculation of (13) in the holomorphic gauge $A_{\bar{z}} = 0$. However, instead of the trivial insertions of q^ρ factors one now needs to insert sophisticated Drinfeld associators [31]. In the holomorphic gauge the theory reduces to the Kontsevich integral of knot [32]. In this case, one can construct the representation of the braid group B_n in $R^{\otimes n}$ through the Drinfeld associators as follows: for the element $g_k \in B_n$ one has:

$$g_k \rightarrow \Psi_k R_k \Psi_k^{-1}, \quad R_i = \underbrace{1 \otimes 1 \otimes \dots \otimes 1}_{1 \dots i-1} \otimes R \otimes \underbrace{1 \otimes \dots \otimes 1}_{i+2 \dots n}, \quad \text{with } R = q^{T^a \otimes T^a} \quad (17)$$

$\Psi_k = \Phi_k \otimes 1^{\otimes (n-k)}$, where Φ_k is the k -th Drinfeld associator. Then, again, if $b_K \in B_n$ is the braid representing some knot K , for the quantum invariant (13) one obtains

$$\langle K \rangle_R = \text{qtr}_R(1)^n q^{-w(b_K) \Omega_2(R)} \text{qtr}_R(b_K) \quad (18)$$

Again, the classical R -matrices $R = q^{T^a \otimes T^a}$ stand just at the intersection points of the K projection on a plane. However, instead of the simple enhancement factors q^ρ , one now inserts in between the quantum associators [31], acting on k lines simultaneously. The Drinfeld associators are solutions to the Knizhnik-Zamolodchikov equations [33] in the WZNW conformal theory [20]. In particular, the k -th associator describes the monodromy of $k + 2$ -point function in the WZNW model, and the Kontsevich integral (18) is nothing but the trace of monodromy associated with the braid b_K . For further details of this representation see [34].

3.3 Representation through Vassiliev invariants and Kontsevich integrals

In CS theory perturbative expansion, the dependencies of the Wilson average $\langle K \rangle_R$ on the knot K and on the group structure G, R are nicely separated:

$$\langle K \rangle_R = \dim_q(R) \prod_{m=0}^{\infty} \prod_{n=1}^{d_m} \exp\left(\hbar^m \alpha_{m,n}(K) r_{m,n}(R)\right) \quad (19)$$

where $\dim_q(R)$ is the quantum dimension of representation:

$$\dim_q(R) = \frac{q^{\frac{N}{2}} - q^{-\frac{N}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}$$

The quantities $\alpha_{m,n}(K)$ are the primary Vassiliev invariants [35], which are rational(!) numbers, naturally represented either as modifications of the Gauss linking integrals in the Lorentz gauge $\partial_\mu A_\mu = 0$ [36], or as the Kontsevich integrals [32] in the holomorphic gauge $A_{\bar{z}} = 0$, or through the writhe numbers [37] in the temporal gauge $A_0 = 0$.

The group factors $r_{m,n}$ are the eigenvalues of operators from the cut-and-join algebra [38] on the $GL(\infty)$ characters χ_R , and form a basis of the *multiplicative independent* Casimir eigenvalues of order m . In every order m one has d_m independent Casimirs, the first several values are:

m	1	2	3	4	5	6
d_m	0	1	1	2	3	5

(20)

$r_{kj}(R)$ can be also understood as eigenvalues of operators from the cut-and-join algebra [38] on the characters χ_R , and are certain non-linear combinations of the Casimir eigenvalues. The Casimir functions $r_{m,n}$ are polynomials of the rank and the weight of the representations, the following table lists the first four Casimirs for $SU(n)$, $SO(n)$ in the fundamental representation and for $SU(2)$ in the spin J representation:

	$r_{2,1}$	$r_{3,1}$	$r_{4,1}$	$r_{4,2}$
$SU(n)$ fund.	$-\frac{n^2-1}{4}$	$-\frac{n(n^2-1)}{8}$	$-\frac{n^2(n^2-1)}{16}$	$-\frac{(n^2+2)(n^2-1)}{16}$
$SO(n)$ fund.	$-\frac{(n-2)(n-1)}{16}$	$-\frac{(n-2)^2(n-1)}{64}$	$-\frac{(n-2)^3(n-1)}{256}$	$\frac{(n-1)(n-2)(n^2-5n+10)}{256}$
$SU(2), J$	$-J(J+1)$	$-J(J+1)$	$-J(J+1)$	$2J^2(J+1)^2$

(21)

Our choice of the basis of $r_{m,n}$ is shown in Fig.2 in the form of the chord diagrams: the circle stands for the trace over representation R , and the trivalent vertices depict the structure constants f^{abc} of the algebra. This notation is clear from the two examples:

$$r_{2,1} = \frac{1}{\dim_R} f^{abc} \text{tr}_R(T^a T^b T^c), \quad r_{3,1} = \frac{1}{\dim_R} f^{abe} f^{ec,d} \text{tr}_R(T^a T^b T^d T^c)$$

In the case of $G = SL(2)$ the relevant representations R are labeled by the value of spin J , and $r_{kj}(J)$ are polynomials of J of degree k . The series in the exponential (19) can be re-expanded in new variables $\hbar$ and $N = 2J + 1 = u/\hbar$:

$$F(K, \hbar, u) = \log \langle K \rangle_R - \log N = \sum_{k=0}^{\infty} \hbar^k \sum_{j=1}^{d_k} \alpha_{kj}(K) r_{kj}(N/\hbar) = \sum_k \hbar^k F_k(K, u) \quad (22)$$




Figure 2: A basis of independent Casimir function up to degree 6.

These $F_k(K, u)$ are infinite combinations of Vassiliev invariants of all orders. The $K - J$ separation gets obscure after such a re-expansion, and all the information about the gauge group ($r_{m,n}$) and the knot ($\alpha_{m,n}$) gets mixed. Instead this expansion is the knot theory counterpart of the genus ($1/N$) expansion in matrix models, and $u = \hbar N$ plays role of the t'Hooft's coupling.

Remarkably, for some knots (called *hyperbolic*) $F(K)$ behaves as $V(K, u)\hbar^{-1}$ at small $\hbar$ [39], this is despite all terms in the series have positive powers of $\hbar$! This is typical for genus expansion series: contributions of every particular genus are all convergent series, but the sum over genera diverges: it can be studied with the help of the Pade summation methods, see [40, 41] for a recent description within the matrix model context. The value of the coefficient $V(K, u = 1)$ at $u = 1$ coincides with the volume of S^3/K in the uniquely defined hyperbolic metric [42], the situation is far more interesting for non-hyperbolic knots (*toric* or *satellite*), see the plots in appendix C.

3.4 Representation through quantum dilogarithm

In [26] the polynomial invariants (13) for the gauge group $SU(2)$ with spin J were systematically interpreted as generalized q -hypergeometric functions, such representations were widely used before in particular examples. This means that these Wilson averages can be represented in the form of finite sums:

$$\langle K \rangle_J = \dim_q(R) \sum_{k_1, k_2, \dots, k_n} \frac{(q, N)_{k_1} \dots (q, N)_{i_1}}{(q, N)_{k_{i_2}} \dots (q, N)_{k_n}} q^{p_2(k_1, \dots, k_n)} \quad (23)$$

Here $N = 2J + 1$, $(q, N)_k$ is the q -Pochhammer symbol:

$$(q, N)_k = \prod_{i=1}^k \left(q^{\frac{N-i}{2}} - q^{-\frac{N-i}{2}} \right) \quad (24)$$

and $p_2(k_1, \dots, k_n)$ is a certain quadratic function. The numbers of k_i in the sum (23), the quadratic polynomial p_2 , the arrangement of the Pochhammer symbols are determined by the knot K and the limits of summation by J . The existence of such a representation for knots directly follows from the AJ -conjecture for the colored

Jones polynomials [43], which states that $\langle K \rangle_J$ is a solution of certain hypergeometric difference equation, see also s.3.6 below. At the same time, the Pochhammer symbols can be expressed through the ratio of the quantum dilogarithms:

$$(q, N)_k = (-i)^k \frac{s \left(i\epsilon_2(N-1-k) + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)}{s \left(i\epsilon_2(N-1) + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)} \quad (25)$$

and the expression (23) can be schematically rewritten as

$$\langle K \rangle_J = \dim_q(R) \sum_{k_1, k_2, \dots, k_n} \frac{s_{k_1} \dots s_{k_{i_1}}}{s_{k_{i_2}} \dots s_{k_n}} q^{p_2(k_1, \dots, k_n)} \quad (26)$$

This provides a discrete version of the Hikami invariants.

3.5 Hikami formalism in Chern-Simons theory

The idea of Hikami formalism is to calculate the CS partition functions, making use of triangulation of the $3d$ manifold M , i.e. by decomposing it into elementary symplexes, tetrahedra. Each tetrahedron has four faces, one can choose two and call them white, then the other two will be black, and one glues black faces to the white ones only. With each site one associates a number p with the white faces, and p^* with the black ones. When two sites are identified, the two numbers are identified, $p_i = p_j^*$. Finally, with each tetrahedron one associates a function $G(p_1^*, p_2^* | p_1, p_2)$, and integrates over all p -variables on identified faces. In this way, with each triangulation of S^3 one associates a multiple integral over all p -variables, one per each 2-face of triangulation:

$$H(K|u) = \int \prod_{\text{simplices}} dp_{i_1} dp_{i_2} dp_{i_1} dp_{i_2} G(p_{i_1}^*, p_{i_2}^* | p_{i_1}, p_{i_2}) \prod_{2\text{-faces}} \delta_{p_{k_m}^* - p_{k_n}} \prod_{1\text{-cycles}} \delta \left(\sum p_j - u \right) \quad (27)$$

At the same time, geometrically, one can associate with the system of the glued tetrahedron a knot K , moreover, the u -variables, which are associated with the 1-cycles, can be interpreted as the $U(1)$ -monodromies around the knot K . Most important, the functions G were found in [14] to be

$$G(p_{i_1}^*, p_{i_2}^* | p_{i_1}, p_{i_2}) = \delta(p_1 + p_2 - p_1^*) s(p_2^* - p_2 - i\pi + \hbar | \epsilon_1, \epsilon_2) e^{p_1(p_2^* - p_2)/2\hbar + \hbar^2/6} \quad (28)$$

and the Hikami integral $H(K|u)$ for the given K has exactly the same form as the Wilson average $\langle K \rangle_R$ from s.3.4 in the $(\hbar, u)$ variables, only the sum is substituted by the integral, see examples in Appendix B below. The transition from sums to integrals is associated with the switching from compact to non-compact groups.

3.6 Spectral curves and topological recursion

At least for $G = SL(2)$ the average $\langle K \rangle_R$ is annihilated by a K -dependent difference operator, i.e. satisfies a recurrent relation in spin J , this statement is sometime called the AJ-conjecture [43] (for the first examples beyond $SL(2)$ see [44]). In the variables $u = N\hbar$, where $N = D_{2J+1} = 2J + 1$ one has:

$$\mathcal{A}(e^{\hbar\partial_u}, e^u) \langle K \rangle_R = 0 \quad (29)$$

In the $\hbar = 0$ limit this operator turns into a function (polynomial), and the difference equation into an algebraic one,

$$\Sigma(K) : \quad \mathcal{A}(w, \lambda) = 0 \quad (30)$$

defining the spectral curve $\Sigma(K)$. One can further define the Seiberg-Witten differential

$$dS = \log w d \log \lambda \quad (31)$$

In the typical examples of 4_1 and $m009$ knots, this is a quadratic equation in w (note that an additional $U(1)$ factor $w - 1$ splits away decreasing degree of the equations by one):

$$\Sigma(K) : \quad \mathcal{A}(w, \lambda) = w + \frac{1}{w} = 2f \left(\lambda + \frac{1}{\lambda} \right), \quad w_{\pm}(\lambda) = f \pm \sqrt{f^2 - 1} \quad (32)$$

and

$$dS = \left(\log w_+(\lambda) - \log w_-(\lambda) \right) d \log \lambda = \log \frac{f + \sqrt{f^2 - 1}}{f - \sqrt{f^2 - 1}} d \log \lambda \quad (33)$$

The free energy (22) $F = \log \langle K \rangle_R$ is actually an exact Seiberg-Witten prepotential, reconstructed in all orders of genus expansion in $\hbar$ from this data (Σ_K, dS) with the help of the AMM/EO topological recursion [27]. See [9] for detailed examples of this reconstruction. One may wonder, *what* is the corresponding SW theory, – and one easily recognizes in (32) and (33) the formulas from [45], describing the $5d$ version of the classical SW theory in terms of the relativistic Toda integrable system. Then (29) should be a Baxter equation, which is now known [46, 47] to describe the NS deformation of the classical SW theory (or, what is the same, the NS limit of the full SW theory). This observation implies a new kind of AGT duality, to be discussed in s.5 below. When $\mathcal{A}(w, \lambda) = 0$ is not reduced to a quadratic equation in w , the analysis is more complicated.

4 A $3d$ AGT relation

Suggestion of [1] is somehow to identify the modular kernels $M(a, a')$, associated with modular transformations of the punctured Riemann surface $S(q) \rightarrow S(q')$, with Chern-Simons cobordism amplitudes on the $3d$ space, which interpolates between $S(q)$ and $S(q')$. In particular, the trace of $M(a, a')$ should be compared with the knot invariants, where the relevant knot K is formed by the closed trajectories of punctures, while external momenta are associated with monodromies around the knot components. Of course, K depends on the choice of the modular transformation. Indeed, there is a striking similarity between both types of quantities, (38) and (39) below. Both are multiple integrals of products of the quantum-dilogarithm functions $s(\dots | \epsilon_1, \epsilon_2)$, and there is a natural identification of parameters:

$$2\pi i \hbar = \log q = \frac{2\pi i}{k + C_G} \stackrel{[1]}{=} 2\pi i b^2 = \frac{2\pi i \epsilon_2}{\epsilon_1} \quad (34)$$

However, there are also differences: in the number of s -functions, in their arguments and in the integration contours. Perhaps, the most striking difference is that the integration variables obey "conservation laws" in expressions for knot invariants, but they do not do so in the trace of monodromy matrix. To see this, one can use the property

$$s(z) = 1/s(-z) \quad (35)$$

to put all the s -functions in the numerator, so that all the integrals acquire the "canonical" form:

$$\prod_i \int_{C_i} dp_i \prod_m s(A_{mi} p_i + B_m | \epsilon_1, \epsilon_2) \exp(C_{ij} p_i p_j + D_i p_i + E) \quad (36)$$

Then

$$\sum_m A_{mi} = 0 \quad (37)$$

in (12), but this is not the case for knot invariants, at least for some i , i.e. for some of the integration variables. The simplest example of such a discrepancy is between

$$T_{\pm}(z) = \int \frac{s(z + \tilde{\alpha})}{s(z - \tilde{\alpha})} e^{\pm i\pi z^2} \quad (38)$$

which are constituents in the expression for the toric modular trace, and

$$\langle 4 \rangle_1 \sim \int s(z + u) s(z - u) e^{\frac{6\pi i u z}{\epsilon_1 \epsilon_2}} dz \quad (39)$$

for the Wilson/Hikami average, associated with the $3d$ AGT-related 4_1 knot. Thus, explicit formulation of the $3d$ AGT hypothesis is clearly very near, but still remains to be found.

5 A route to alternative AGT relation

A seeming failure of the 3+3 AGT relation can attract some new attention to possible alternatives: one actually expects a lot of different AGT like relations to exist. An obvious possibility is to look for an extension of $2+4$ to $3+5$, and try to relate quantities in the $3d$ Chern-Simons theory with those in the $5d$ SYM. This is of course a far more straightforward exercise. Equations in section 3.6 above describe knot invariants, but it is easy to recognize in (32) the spectral curve for the relativistic Toda system, and in (33) the corresponding Seiberg-Witten differential [45]. Thus, these equations establish a clear link between $3d$ Chern-Simons and the $5d$ version of Seiberg-Witten theory described as a straightforward q -deformation of the ordinary $4d$ SW theory.

Within this identification, one should associate the corresponding difference equation (29) with the Baxter equation for the same system. The Baxter equations are known [46] to arise in the Nekrasov-Shatashvili limit [48] and, therefore, one obtains a new AGT like relation:

$$3d \text{ Chern - Simons} \xleftrightarrow{\text{new AGT}} \text{NS limit of } 5d \text{ LMNS prepotential} \xleftrightarrow{\text{ordinary AGT [49]}} q - \text{Virasoro conformal blocks}$$

More concretely, in accordance with [46], one associates with the solution to the Baxter equation, i.e. with $\langle K \rangle_R$, the SW differential, its monodromies around the A - and B -cycles on the spectral curve (32) giving rise to the $5d$ Nekrasov functions in the NS limit via the SW equations.

There are a few interesting points to be mentioned already at this stage. The dilogarithm formulae for the knot invariants from ss.3.4 and 3.5 provide integral representations for solutions of equations (29). They look very similar to solutions [50] for the *open* quantum relativistic Toda chain system, and such solutions are *not* available for the *closed* chain, but (32) is of this latter kind! The thing is that, first, (29) defines a Baxter equation at some special point of the moduli space: for special values of "energies", thus, the fact that solutions are unavailable in such a form at a generic point does not forbid them to exist at some special point. Second, while the classical equations (32) are clearly of the relativistic Toda type, their quantization is not unique, and (29) is not the standard version of the Baxter equation, considered in the literature. One more important point is that when dealing with difference (rather than differential) equations, one obtains infinitely more solutions. To fix this ambiguity one imposes two difference equations where one would impose only one differential. And this pair of equations is typically related by the symmetry $\epsilon_1 \leftrightarrow \epsilon_2$, which is explicitly broken by all our construction of knot invariants. In particular, it is explicitly broken in the basic AGT identification (34).

An interesting question is what should substitute the knot invariants when this new AGT duality is lifted to the entire LMNS deformation of $5d$ theory, beyond the Nekrasov-Shatashvili limit.

6 Conclusion

After discovery of the AGT relation [13], which embeds the $2d$ conformal theory into the general context of Seiberg-Witten and integrability theory [51], a hunt has immediately started for its extension, which would do the same with the $3d$ Chern-Simons theory. The goal of this letter is to switch the discussion of the $3d$ AGT relations from the qualitative to a quantitative mode. This is made possible by the extensive progress in the theory of knot invariants, which is briefly reviewed in s.3 of the present letter. (Actually, some derivations from CS theory itself are still lacking, but this is mostly because of the insufficient attention to these important problems within the QFT community.) Given the existing set of explicit formulas, one can easily test various suggestions. In this way we point out some problems with the suggestion by [1] to AGT-relate the knot invariants with the modular kernels. Instead, we demonstrate that these invariants are AGT-related to the $5d$ SYM theory, this is a more straightforward and less intriguing option, still it also deserves the attention.

Acknowledgements

We acknowledge useful consultations from P.Dunin-Barkovsky, S.Kharchev and A.Sleptsov. Our work is partly supported by Ministry of Education and Science of the Russian Federation under contract 02.740.11.5194, by RFBR grants 10-01-00536 and by joint grants 11-02-90453-Ukr, 09-02-93105-CNRS, 09-02-91005-ANF, 10-02-92109-Yaf-a, 11-01-92612-Royal Society.

Appendix A. Dilogarithm properties

A big problem with discussions of 3d AGT relations is the lack of common notations: people coming from different fields use definitions of the same quantities, which differ by all kinds of rescalings. The purpose of this Appendix is to list at least some relations between various definitions of quantum dilogarithms used in the literature: $S(z|\epsilon_1, \epsilon_2)$, $S_b(z)$, $\Phi_h(z)$, $e_b(z)$. We also demonstrate the trick needed to take the massless limit $\alpha \rightarrow 0$ of the modular kernel $M(a, a')$, i.e. to derive eq.(8).

A1. Various dilogarithms

The function $s(z)$ is the "quantum dilogarithm" [18], the ratio of two digamma-functions [17],

$$s(z|\epsilon_1, \epsilon_2) = \prod_{m, n \geq 0} \frac{(m + \frac{1}{2})\epsilon_1 + (n + \frac{1}{2})\epsilon_2 - iz}{(m + \frac{1}{2})\epsilon_1 + (n + \frac{1}{2})\epsilon_2 + iz} = \frac{\Gamma_2(\epsilon/2 + iz|\epsilon_1, \epsilon_2)}{\Gamma_2(\epsilon/2 - iz|\epsilon_1, \epsilon_2)} \quad (40)$$

It enjoys a set of periodicity properties

$$s\left(z - \frac{i\epsilon_2}{2}|\epsilon_1, \epsilon_2\right) = 2 \cosh\left(\frac{\pi z}{\epsilon_1}\right) s\left(z + \frac{i\epsilon_2}{2}|\epsilon_1, \epsilon_2\right) \quad (41)$$

$$s\left(z - \frac{i\epsilon_1}{2}|\epsilon_1, \epsilon_2\right) = 2 \cosh\left(\frac{\pi z}{\epsilon_2}\right) s\left(z + \frac{i\epsilon_1}{2}|\epsilon_1, \epsilon_2\right) \quad (42)$$

$$s\left(z - \frac{i\epsilon}{2}|\epsilon_1, \epsilon_2\right) = 4 \sinh\left(\frac{\pi z}{\epsilon_1}\right) \sinh\left(\frac{\pi z}{\epsilon_2}\right) s\left(z + \frac{i\epsilon}{2}|\epsilon_1, \epsilon_2\right) \quad (43)$$

This definition of the quantum dilogarithm admits the integral representation

$$i \log s(z|\epsilon_1, \epsilon_2) = \int_0^\infty \frac{dw}{w} \left\{ \frac{\sin(2zw)}{2 \sinh(\epsilon_1 w) \sinh(\epsilon_2 w)} - \frac{z}{\epsilon_1 \epsilon_2 w} \right\} \quad (44)$$

which can be used to derive the asymptotic formulae

$$i \log s(z|\epsilon_1, \epsilon_2) = \frac{\pi z^2}{2\epsilon_1 \epsilon_2} - \frac{\pi}{24} \frac{2\epsilon_1 \epsilon_2 - \epsilon^2}{\epsilon_1 \epsilon_2} + i \sum_{n=0}^\infty \frac{B_n(1/2)}{n!} \left(2\pi i \frac{\epsilon_2}{\epsilon_1}\right)^{n-1} \text{Li}_{2-n}\left(-e^{\frac{2\pi z}{\epsilon_1}}\right) \quad (45)$$

The following resummation expansion is also of use

$$i \log s(z_0 + z|\epsilon_1, \epsilon_2) = \frac{\pi(z_0 + z)^2}{2\epsilon_1 \epsilon_2} - \frac{\pi}{24} \frac{2\epsilon_1 \epsilon_2 - \epsilon^2}{\epsilon_1 \epsilon_2} + \sum_{k=-1}^\infty \sum_{j=0}^\infty \frac{i^{k+1} B_{k+1}(1/2)}{(k+1)! j!} (2\pi)^{k+j} \frac{\epsilon_2^k}{\epsilon_1^{k+j}} \text{Li}_{1-j-k}\left(-e^{\frac{2\pi z_0}{\epsilon_1}}\right) z^j \quad (46)$$

The quantum dilogarithm is symmetric w.r.t. ϵ_1 and ϵ_2 , however, we have explicitly chosen ϵ_2 to be small here, this expansion playing a crucial role in the NS limit.

This definition of the quantum dilogarithm is convenient for applications in the context of AGT conjecture, where parameters $\epsilon_{1,2}$ are explicitly specified. Though in the literature there is another definition which differs by rescaling

$$S_b(z) = \exp\left(\frac{1}{i} \int_0^\infty \frac{dw}{w} \left(\frac{\sin 2zw}{2 \sinh bw \sinh b^{-1}w} - \frac{z}{w}\right)\right) \quad (47)$$

so that

$$\boxed{s(z|\epsilon_1, \epsilon_2) = S_{\sqrt{\frac{\epsilon_1}{\epsilon_2}}}\left(\frac{z}{\sqrt{\epsilon_1 \epsilon_2}}\right)} \quad (48)$$

A2. Modular kernel

In this section we present a standard trick to compute the modular kernel (5) in the simple limit of zero external dimension $\tilde{\alpha} = i\frac{\epsilon}{2}$. Considering the modular kernel in this limit naively, one derives from eq.(43) that

$$M = 2^{3/2} \int dr \frac{e^{4\pi i a' r}}{16 \sinh\left(\frac{\pi(a+r)}{\epsilon_1}\right) \sinh\left(\frac{\pi(a+r)}{\epsilon_2}\right) \sinh\left(\frac{\pi(a-r)}{\epsilon_1}\right) \sinh\left(\frac{\pi(a-r)}{\epsilon_2}\right)} \quad (49)$$

The denominator in the integrand has double poles, which were glued together in the limit $\tilde{\alpha} \rightarrow i\frac{\epsilon}{2}$, and the chosen integration contour is pinched between them. Hence, one has to take the limit more carefully:

$$\begin{aligned} \mathcal{M}(a, a'|0) &= \oint_{r=-a} dr \frac{s(a+r+i\epsilon/2+i\lambda)}{s(a+r-i\epsilon/2-i\lambda)} \frac{e^{4\pi i a' r}}{4 \sinh\left(\frac{\pi(a-r)}{\epsilon_1}\right) \sinh\left(\frac{\pi(a-r)}{\epsilon_2}\right)} + \\ &+ \oint_{r=a} dr \frac{s(a-r+i\epsilon/2+i\lambda)}{s(a-r-i\epsilon/2-i\lambda)} \frac{e^{4\pi i a' r}}{4 \sinh\left(\frac{\pi(a+r)}{\epsilon_1}\right) \sinh\left(\frac{\pi(a+r)}{\epsilon_2}\right)} \end{aligned} \quad (50)$$

Since

$$\begin{aligned} \frac{s(a+r+i\epsilon/2+i\lambda)}{s(a+r-i\epsilon/2-i\lambda)} &= \prod_{m,n \geq 0} \frac{(m+1/2)\epsilon_1 + (n+1/2)\epsilon_2 - i(a+r+i\epsilon/2+i\lambda)}{(m+1/2)\epsilon_1 + (n+1/2)\epsilon_2 + i(a+r+i\epsilon/2+i\lambda)} \times \\ &\times \frac{(m+1/2)\epsilon_1 + (n+1/2)\epsilon_2 + i(a+r-i\epsilon/2-i\lambda)}{(m+1/2)\epsilon_1 + (n+1/2)\epsilon_2 - i(a+r-i\epsilon/2-i\lambda)} = \\ &= \prod_{m,n \geq 0} \frac{(m+1)\epsilon_1 + (n+1)\epsilon_2 - i(a+r+i\lambda)}{m\epsilon_1 + n\epsilon_2 + i(a+r+i\lambda)} \frac{(m+1)\epsilon_1 + (n+1)\epsilon_2 + i(a+r-i\lambda)}{m\epsilon_1 + n\epsilon_2 - i(a+r-i\lambda)} \sim \\ &\sim_{m,n=0} \frac{1}{(a+r)^2 - \lambda^2} \sim_{\lambda \rightarrow 0} \delta(a+r) \end{aligned} \quad (51)$$

one finally obtains

$$\mathcal{M}(a, a'|0) \rightarrow \frac{\sqrt{2} \cos\left(4\pi i \frac{aa'}{\epsilon_1 \epsilon_2}\right)}{\mu'(a)} \quad (52)$$

A3. Chern-Simons average

Standard quantities arising in the Chern-Simons theory often use another definition of the dilogarithm. For instance, the 4_1 -knot average is usually written as a function of the coupling constant $\hbar$ and the knot monodromy parameter u as (note that $\hbar$ in our formulae differs from h in [14] by a factor of 2)

$$\langle 4_1 \rangle = H(u, \hbar) = \frac{1}{\sqrt{\pi \hbar}} \int dp \frac{\Phi_{\hbar}(p + i\pi + \hbar/2)}{\Phi_{\hbar}(-2u - p - i\pi - \hbar/2)} e^{-\frac{4}{\hbar} u(u+p) - u} \quad (53)$$

where

$$\Phi_{\hbar}(z) = \Phi\left(\frac{z}{\pi i \hbar} \middle| \frac{\hbar}{2\pi i}\right), \quad \Phi(z|\tau) = \exp\left(\frac{1}{4} \int \frac{dw}{w} \frac{e^{2xz}}{\sinh w \sinh \tau w}\right) \quad (54)$$

In the previously discussed context of CFT one can encounter a similar function though with rescaled parameters

$$e_b(z) = \exp\left(\frac{1}{4} \int \frac{dw}{w} \frac{e^{-2izw}}{\sinh bw \sinh b^{-1}w}\right) = \Phi(-ibz|b^2) \quad (55)$$

so that

$$\langle 4_1 \rangle = \frac{1}{\sqrt{2\pi(\pi i b^2)}} \int dp \frac{e_b\left(\frac{1}{2\pi b}(p + i\pi + \pi i b^2)\right)}{e_b\left(\frac{1}{2\pi b}(-2u - p - i\pi - \pi i b^2)\right)} e^{-\frac{2}{\pi i b^2} u(u+p) - u} \quad (56)$$

Introducing new variables $p = 2\pi b z$, $u = 2\pi b u'$, $Q = b + b^{-1}$, one obtains

$$\langle 4_1 \rangle = \frac{2\pi b}{\sqrt{2\pi(\pi i b^2)}} \int dz \frac{e_b(z + iQ/2)}{e_b(-2u' - z - iQ/2)} e^{8\pi i u'(u'+z) - 2\pi b u'} \quad (57)$$

Notice that, though similar, the functions e_b and S_b are slightly different, in particular, they differ by a multiplier:

$$e_b(z) = e^{\frac{\pi i z^2}{2}} e^{-\frac{i\pi(2-Q^2)}{24}} S_b(z) \quad (58)$$

Therefore,

$$\begin{aligned} \langle 4_1 \rangle &= \sqrt{-2i} \int dz \frac{S_b(z + iQ/2)}{S_b(-2u' - z - iQ/2)} e^{6\pi i u'(z+u') - \pi u'(b-b^{-1})} = \\ &=_{z \rightarrow z-u'-iQ/2} \sqrt{-2i} \int dz S_b(z - u') S_b(z + u') e^{6\pi i u' z + 2\pi u'/b + 4\pi u'b} \end{aligned} \quad (59)$$

At last, the same expression in terms of $s(z|\epsilon_1, \epsilon_2)$ -dilogarithms is

$$H\left(\frac{2\pi u}{\sqrt{\epsilon_1 \epsilon_2}}, \pi i \frac{\epsilon_2}{\epsilon_1}\right) = \sqrt{\frac{-2i}{\epsilon_1 \epsilon_2}} \int dz s(z - u|\epsilon_1, \epsilon_2) s(z + u|\epsilon_1, \epsilon_2) e^{\frac{6\pi i u z}{\epsilon_1 \epsilon_2} + \frac{2\pi u}{\epsilon_2} + \frac{4\pi u}{\epsilon_1}} \quad (60)$$

A4. Pochhammer symbols as dilogarithm ratios

Consider the Pochhammer symbols

$$\begin{aligned} (q, N)_k &= \prod_{j=1}^k \left(q^{\frac{N-j}{2}} - q^{-\frac{N-j}{2}} \right) = 2^k \prod_{j=1}^k \sinh(\pi i \hbar(N-j)), \\ (q, N)_k^* &= \prod_{j=1}^k \left(q^{\frac{N+j}{2}} - q^{-\frac{N+j}{2}} \right) = 2^k \prod_{j=1}^k \sinh(\pi i \hbar(N+j)) \end{aligned} \quad (61)$$

Using the periodicity conditions (43) one can rewrite the sine products through

$$\sinh\left(\frac{\pi i \epsilon_2 z}{\epsilon_1}\right) = -\frac{i}{2} \frac{s\left(i\epsilon_2 z + \frac{i}{2}(\epsilon_1 - \epsilon_2)\right)}{s\left(i\epsilon_2 z + \frac{i}{2}(\epsilon_1 + \epsilon_2)\right)} \quad (62)$$

i.e. the Pochhammer symbols are

$$(q, N)_k = (-i)^k \prod_{j=1}^k \frac{s(i\epsilon_2(N-1-j) + i\epsilon/2)}{s(i\epsilon_2(N-j) + i\epsilon/2)} = (-i)^k \frac{s(i\epsilon_2(N-1-k) + i\epsilon/2)}{s(i\epsilon_2(N-1) + i\epsilon/2)} \quad (63)$$

and, correspondingly,

$$(q, N)_k^* = (-i)^k \frac{s(i\epsilon_2(N) + i\epsilon/2)}{s(i\epsilon_2(N+k) + i\epsilon/2)} \quad (64)$$

Note that $\langle 4_1 \rangle$ is expressed through $(q, N/\sqrt{\hbar})_{k/\sqrt{\hbar}}$, which celebrates the symmetry $\hbar \rightarrow \hbar^{-1}$:

$$\langle 4_1 \rangle = \frac{i}{\cosh(i\epsilon_2(N-1/2) + i\epsilon/2)} \sum_{k \in \mathbb{Z} + \frac{1}{2}} (-1)^k \frac{s(i\epsilon_2(N-1/2-k) + i\epsilon/2)}{s(i\epsilon_2(N-1/2+k) + i\epsilon/2)} \quad (65)$$

In the limit $\epsilon_2 \rightarrow 0$, $N \rightarrow \infty$, $i\epsilon_2(N-1/2) + i\epsilon/2 = \tilde{u}$ the sum can be substituted by the integral to give

$$\langle 4_1 \rangle = -\frac{1}{\epsilon_2 \cosh \tilde{u}} \int dz e^{\pi z/\epsilon_2} s(z - \tilde{u}) s(z + \tilde{u}) \quad (66)$$

which is quite similar to the Hikami formula, though the correct exponential factor is not restored in the integrand.

Appendix B. Examples of knots invariants

In this section we provide a few simplest examples of knots and all the associated elements of their description, outlined in section 3: for each knot K we list the corresponding annihilating operator polynomial $\mathcal{A}$, the spectral curve, the braid-group element responsible for the quantum- R -matrix description of $\langle K \rangle_R$, the relevant combination of classical R -matrix and Drinfeld associator, the expansion of $F = \log \langle K \rangle_R$ in the Vassiliev invariants, the representation of $\langle K \rangle_R$ and the Hikami integral $H(K|u)$ through relevant combinations of s -functions.

B1. Unknot

- **Quantum R-matrix representation**

The simplest braid representation of the unknot U_0 is the closure of single string, i.e. the closure of the only element of group B_1 :

$$b_{U_0} = 1 \in B_1 \quad (67)$$

Thus, the value of polynomial invariant is given by the following character:

$$\langle U_0 \rangle = \text{qtr}_R(1) = \text{tr}_R(q^\rho) = \chi_R(z_i = q^{N-2i+1})$$

where χ_R is the character of representation R (the corresponding Schur polynomial).

- **Representation through Drinfeld associator**

Formula (18) for the unknot gives vev of unknot in the form of "hump" trace of the first Drinfeld associator Φ_3 . Relation between the two is non-trivial: the $\langle \text{hump} \rangle$ is rather inverse of $\langle U_0 \rangle$. The associator is a tensor with six indices, and the hump trace is defined as the following contraction:

$$\langle \text{hump} \rangle = \sum_{i,k,m=1}^{\dim_R} \left(\Phi_3 \right)_{k,m,m}^{i,i,k} \quad (68)$$

In the case of fundamental representation of $SU(N)$ this sum was computed explicitly, and the result can be represented as a particular value of the hypergeometric function (see [34] for some details):

$$\langle \text{hump} \rangle = \frac{N}{F\left([(N-1)\hbar, (N+1)\hbar], [1+N\hbar], 1\right)} = \frac{N(q^{\frac{1}{2}} - q^{-\frac{1}{2}})}{q^{\frac{N}{2}} - q^{-\frac{N}{2}}} = \frac{N}{[N]_q}$$

- **Representation through Vassiliev invariant**

The first Vassiliev invariants of unknot are:

$\alpha_{2,1}$	$\alpha_{3,1}$	$\alpha_{4,1}$	$\alpha_{4,2}$	$\alpha_{5,1}$	$\alpha_{5,2}$	$\alpha_{5,3}$
$-\frac{2}{3}$	0	$\frac{2}{45}$	$-\frac{2}{45}$	0	0	0

(69)

For example the usage of (21) and table (69) for the group $SU(2)$ with spin J gives¹:

$$\langle U_0 \rangle_J = N \exp \left(h^2 \frac{2}{3} J(J+1) - h^4 \frac{2}{45} (J+1)(2J^2 + 2J + 1)J + \dots \right) = \frac{q^{\frac{N}{2}} - q^{-\frac{N}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}} = [N]_q \quad (70)$$

with $N = 2J + 1$.

- **$\mathcal{A}$ -polynomial**

The colored Jones polynomial of unknot

$$K_N(U_0|q) = \langle U_0 \rangle_J = \frac{q^{J+\frac{1}{2}} - q^{-(J+\frac{1}{2})}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}} = \frac{q^{\frac{N}{2}} - q^{-\frac{N}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}} \quad (71)$$

is a character (68), and, therefore, it satisfies the quantum Laplace equation on the Cartan lattice:

$$K_{N+1} - [2]_q K_N + K_{N-1} = 0 \quad (72)$$

Therefore, the $\mathcal{A}$ -polynomial for the unknot can be defined as

$$\mathcal{A}(l, m) = (l-1)^2/2, \quad (73)$$

and the quantum $\mathcal{A}$ -polynomial is the q -Laplace operator

$$\hat{\mathcal{A}} = \Delta_q = \hat{l} - [2]_q + \hat{l}^{-1}, \quad \Delta_q K_N = 0 \quad (74)$$

where operators $\hat{l}$ and $\hat{m}$ act on the Jones polynomial as

$$\hat{l}K_N = K_{N+1}, \quad \hat{m}K_N = q^N K_N \quad (75)$$

¹Hereafter, we use the expansions in parameter $h := 2\pi i \hbar$ such that $q = e^h$.

- **Polynomial invariants**

Colored HOMFLY polynomial:

$$\langle U_0 \rangle_Y = \dim_q(Y) = \prod_{i,j \in Y} \frac{q^{\frac{N+j-i}{2}} - q^{-\frac{N+j-i}{2}}}{q^{\frac{h(i,j)}{2}} - q^{-\frac{h(i,j)}{2}}} \quad (76)$$

see [52]. Here Y is the Young diagram corresponding to the representation of $SU(N)$ and $h(i,j)$ is the hook length of a box in Y .

Superpolynomial:

$$P_0(a, q, t) = \frac{a^{\frac{1}{2}} - a^{-\frac{1}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}} \quad (77)$$

Colored superpolynomial: [23, formulae (67)-(68)].

B2. Knot 3_1

- **Representation through quantum R -matrix and Drinfeld associator**

The polynomial invariant for 3_1 can be constructed using formulae (15) and (18). In this case, 3_1 is the closure of the following element:

$$b_{3_1} = g_1^3 \in B_2 \quad (78)$$

In this representation the knot has 3 positively oriented crossings ($w(b) = 3$) and two strings ($n = 2$ in (18)).

- **Representation through Vassiliev invariant**

The first Vassiliev invariants of knot 3_1 are:

$\alpha_{2,1}$	$\alpha_{3,1}$	$\alpha_{4,1}$	$\alpha_{4,2}$	$\alpha_{5,1}$	$\alpha_{5,2}$	$\alpha_{5,3}$
4	-8	$\frac{62}{3}$	$\frac{10}{3}$	$-\frac{176}{3}$	$-\frac{32}{3}$	-8

(79)

For example the usage of (21) for $SU(2)$ gives:

$$\langle 3_1 \rangle = [2J+1]_q \exp \left(-4J(1+J)h^2 + 8J(1+J)h^3 + \frac{2}{3}J(1+J)(-31+10J+10J^2)h^4 + \dots \right) \quad (80)$$

with $h = 2\pi i\hbar$.

- **Representation through quantum dilogarithm**

The colored Jones polynomial for 3_1 with $N = 2J+1$ can be represented in the hypergeometric form:

$$K_N(3_1|q) = [N]_q \sum_{i=0}^{N-1} (-1)^i q^{\frac{i(i+3)}{2}} (q, N)_i (q, N)_i^* \equiv [N]_q J_N(3_1|q) \quad (81)$$

The (63) gives:

$$J_N \sim \sum_{k=0}^{N-1} q^{\frac{k(k+3)}{2}} \frac{s \left(i\epsilon_2(N-1-k) + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)}{s \left(i\epsilon_2(N-1) + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)} \frac{s \left(i\epsilon_2 N + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)}{s \left(i\epsilon_2(N+k) + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)} \quad (82)$$

In the limit of $N \rightarrow \infty$ and $|q| > 1$ this expression turns into

$$J_N \sim q^{\frac{3}{2}N^2 - \frac{3}{4}N} \quad (83)$$

- **$\mathcal{A}$ -polynomial**

The $\mathcal{A}$ -polynomial and the spectral curve for the knot 3_1 have the following form:

$$\mathcal{A}_{3_1}(l, m) = m^3 + l, \quad \Sigma_{3_1} : \{(m, l) \in \mathbb{C}^2 : m^3 + l = 0\} \quad (84)$$

Note that this spectral curve corresponds to the sphere, and to the open relativistic Toda system.

In order to compute the quantum $\mathcal{A}$ -polynomial, let us note that the Jones polynomial for the trefoil (81) satisfies the difference equation

$$J_{N+1} + q^{3N+2} \frac{1 - q^N}{1 - q^{N+1}} J_N = q^N \frac{q^{2N+1} - 1}{q^{N+1} - 1} \quad (85)$$

which can be rewritten as

$$\frac{1}{\hat{B}(\hat{m})} \hat{A}(\hat{l}, m) J_N(q) = 1 \quad (86)$$

with

$$\hat{A} = q\hat{m}^3(\hat{m} - 1) + (q\hat{m} - 1)\hat{l}, \quad \hat{B} = (q\hat{m}^2 - 1)\hat{m} \quad (87)$$

Equivalently, (86) can be rewritten as

$$\sqrt{\frac{q}{\hat{m}^3}} \frac{1 - \hat{m}}{1 - q\hat{m}^2} \left(\hat{l} + q^{\frac{3}{2}} \hat{m}^3 \right) K_N(q) = 1 \quad (88)$$

In the leading order at the large N limit this equation reduces to

$$\left(\hat{l} + q^{\frac{3}{2}} \hat{m}^3 \right) K_N(q) = 0 \quad (89)$$

i.e. the quantum $\mathcal{A}$ -polynomial is²

$$\hat{\mathcal{A}}_{3_1}(\hat{l}, \hat{m}|q) = \hat{l} + q^{\frac{3}{2}} \hat{m}^3 \quad (90)$$

(83) definitely solves (89).

Note that at finite N (88) can be rewritten as

$$\hat{\mathcal{A}}_g J_N(q) \equiv (\hat{l} - 1) \frac{1}{\hat{B}(\hat{m})} \hat{A}(\hat{l}, \hat{m}) J_N(q) = 0 \quad (91)$$

$$\hat{\mathcal{A}}_g = -\frac{1}{q\hat{m}} \frac{1 - q^2\hat{m}}{1 - q^2\hat{m}^2} \hat{l}^2 - \left[\frac{1}{\hat{m}} \frac{1 - q\hat{m}}{1 - q\hat{m}^2} - q^2\hat{m}^2 \frac{1 - q\hat{m}}{1 - q^3\hat{m}} \right] \hat{l} - q^2\hat{m}^2 \frac{1 - \hat{m}}{1 - q^2\hat{m}} \quad (92)$$

- **Polynomial invariants**

Colored HOMFLY polynomial: The colored polynomial is known in this case only for $SU(3)$, see [44, Theorem 1 of the 1st paper].

Superpolynomial (non-colored):

$$P_{3_1}(a, q, t) = P_0(a, q, t) [aq^{-1} + aqt^2 + a^2t^3] \quad (93)$$

Choosing $t = -1$ one obtains from this superpolynomial the HOMFLY polynomial. The specialization of this latter $a = q^N$ corresponds to $SU(N)$ Chern-Simons theory, while $a = q^2$ gives the Jones polynomial and $a = q^0 = 1$ gives the Alexander polynomial.

²In the literature, one often makes the different choice of variables: $q \rightarrow q^2$ and $\hat{m} \rightarrow \hat{m}^2$.

B3. Knot 4_1

- **Representation through quantum R -matrix and Drinfeld associator**

The polynomial invariant for 4_1 can be constructed using formulae (15) and (18). In this case, 4_1 can be represented as the closure of the following element:

$$b_{4_1} = g_2^2 g_1^{-1} g_2 g_1^{-1} \in B_3 \quad (94)$$

In this representation the knot has 3 positively oriented crossings and two negative ones ($w(b) = 1$) and two strings ($n = 3$ in (18)).

- **Representation through Vassiliev invariant**

The first Vassiliev invariants of knot 4_1 are:

$\alpha_{2,1}$	$\alpha_{3,1}$	$\alpha_{4,1}$	$\alpha_{4,2}$	$\alpha_{5,1}$	$\alpha_{5,2}$	$\alpha_{5,3}$
-4	0	$\frac{34}{3}$	$\frac{14}{3}$	0	0	0

(95)

For instance, the usage of (21) for $SU(2)$ gives:

$$\begin{aligned} \langle 4_1 \rangle = [N]_q \exp \left\{ J(J+1)h^2 \left[4 + \frac{2}{3}(14J^2 + 14J - 17)h^2 + \frac{1}{90}(2416J^4 + 4832J^3 - 9212J^2 - 11628J + 8013)h^4 + \right. \right. \\ \left. \left. + \frac{1}{1260}(109552J^6 + 328656J^5 - 973888J^4 - 2495536J^3 + 1783146J^2 + 3085690J - 1645097)h^6 + \dots \right] \right\} \quad (96) \end{aligned}$$

Naively, in the limit, where $u = hN$ is finite, while $N \rightarrow \infty$ (as usual, $N = 2J + 1$) only the terms of degree 0 in N survive, and the series gets the form

$$\lim_{N \rightarrow \infty} \ln \left(\langle 4_1 \rangle \right) = 1 + u^2 + \frac{7}{12}u^4 + \frac{151}{360}u^6 + \dots = \sum_{n=0}^{\infty} \frac{2u^{2n}}{(2n)!} \sum_{k=0}^{\infty} \frac{k^{2n-1}}{\left(\frac{1+\sqrt{5}}{2}\right)^2} = -\ln \left(\left(1 - \frac{m}{\gamma}\right) \left(1 - \frac{1}{\gamma m}\right) \right) \quad (97)$$

where $\gamma = (3 + \sqrt{5})/2$ and $m = e^u$. No $1/\hbar$ -contribution that would correspond to genus $1/2$ in the t'Hooft expansion is seen in this way.

- **Representation through quantum dilogarithm**

The colored Jones polynomial for 4_1 with $N = 2J + 1$ can be represented in the hypergeometric form:

$$K_N(4_1|q) = [N]_q \sum_{i=0}^{N-1} (q, N)_i (q, N)_i^* \equiv [N]_q J_N(4_1|q) \quad (98)$$

Then, (63) gives:

$$K_N \sim \sum_{k=0}^{N-1} (-1)^k \frac{s \left(i\epsilon_2(N-1-k) + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)}{s \left(i\epsilon_2(N-1) + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)} \frac{s \left(i\epsilon_2 N + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)}{s \left(i\epsilon_2(N+k) + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)} \quad (99)$$

- **Hikami Representation**

To proceed from the previous formula (99) to the Hikami integral representation, one needs to go to the double limit $\epsilon_2 \rightarrow 0$, $N \rightarrow \infty$, $i\epsilon_2(N-1/2) + i\epsilon/2 = \tilde{u}$. In this limit, the sum in (99) should be replaced by the integral, and one finally gets

$$K_N = -\frac{1}{\epsilon_2 \cosh \tilde{u}} \int dz e^{\pi z / \epsilon_2} s \left(z - \tilde{u} \middle| \epsilon_1, \epsilon_2 \right) s \left(z + \tilde{u} \middle| \epsilon_1, \epsilon_2 \right) \quad (100)$$

In Hikami's terms, the presence of two dilogarithm functions in the integral (100) shows that the hyperbolic space $S^3 \setminus 4_1$ can be obtained by gluing two tetrahedrons. This representation of $S^4 \setminus 4_1$ is well studied in the literature, see, e.g., [54].

- **$\mathcal{A}$ -polynomial**

The $\mathcal{A}$ -polynomial and the spectral curve for the knot 4_1 have the following form:

$$\mathcal{A}_{4_1}(l, m) = m^2 + l(-1 + m + 2m^2 + m^3 - m^4) + l^2 m^2, \quad \Sigma_{4_1} = \{(m, l) \in \mathbb{C}^2 : \mathcal{A}_{4_1}(l, m) = 0\} \quad (101)$$

In order to calculate the quantum $\mathcal{A}$ -polynomial in this case, we find the difference equation (86) for knot 4_1 with

$$\begin{aligned} \hat{A} &= q^2 \hat{m}^2 (1 - \hat{m})(1 - q^3 \hat{m}^2) - (q\hat{m} + 1)(1 - q\hat{m} - q\hat{m}^2 - q^3 \hat{m}^2 - q^3 \hat{m}^3 + q^4 \hat{m}^4)(1 - q\hat{m})^2 \hat{l} + \\ &\quad + q^2 \hat{m}^2 (1 - q\hat{m}^2)(1 - q^2 \hat{m}) \hat{l}^2, \quad \hat{B} = q\hat{m}(1 - q^3 \hat{m}^2)(1 - q\hat{m}^2)(1 + q\hat{m}) \end{aligned} \quad (102)$$

At the large N limit it reduces to

$$\begin{aligned} &\left[q^{3/2} \hat{m}^2 (1 - q^3 \hat{m}^2) - (1 - q^2 \hat{m}^2)(1 - q\hat{m} - q\hat{m}^2 - q^3 \hat{m}^2 - q^3 \hat{m}^3 + q^4 \hat{m}^4) \hat{l} + \right. \\ &\quad \left. + q^{5/2} \hat{m}^2 (1 - q\hat{m}^2) \hat{l}^2 \right] K_N(q) = 0 \end{aligned} \quad (103)$$

and is solved by formula (100). Thus, the quantum $\mathcal{A}$ -polynomial is

$$\hat{\mathcal{A}}_{4_1}(\hat{l}, \hat{m}|q) = q^{3/2} \hat{m}^2 (1 - q^3 \hat{m}^2) - (1 - q^2 \hat{m}^2)(1 - q\hat{m} - q\hat{m}^2 - q^3 \hat{m}^2 - q^3 \hat{m}^3 + q^4 \hat{m}^4) \hat{l} + q^{5/2} \hat{m}^2 (1 - q\hat{m}^2)$$

At finite N the Jones polynomial satisfies the equation

$$\hat{\mathcal{A}}_g J_N(q) \equiv (\hat{l} - 1) \frac{1}{\hat{B}(\hat{m})} \hat{A}(\hat{l}, \hat{m}) J_N(q) = 0 \quad (104)$$

$$\begin{aligned} \hat{\mathcal{A}}_g &= -\frac{q^2 \hat{m} (1 - q^3 \hat{m})}{(1 + q^2 \hat{m})(1 - q^5 \hat{m}^2)} \hat{l}^3 + \\ &\quad + \frac{1}{q^2} \frac{(1 - q^2 \hat{m})(1 + q\hat{m} - 2q^2 \hat{m} - q^3 \hat{m}^2 + q^4 \hat{m}^2 - q^5 \hat{m}^2 - 2q^6 \hat{m}^3 + q^7 \hat{m}^3 + q^8 \hat{m}^4)}{\hat{m}(1 + q\hat{m})(1 - q^5 \hat{m}^2)} \hat{l}^2 - \\ &\quad - \frac{1}{q} \frac{(1 - q\hat{m})(1 - 2q\hat{m} + q^2 \hat{m} - q\hat{m}^2 + q^2 \hat{m}^2 - q^3 \hat{m}^2 + q^2 \hat{m}^3 - 2q^3 \hat{m}^3 + q^4 \hat{m}^4)}{\hat{m}(1 + q^2 \hat{m})(1 - q\hat{m}^2)} \hat{l} + \frac{q\hat{m}(1 - \hat{m})}{(1 + q\hat{m})(1 - q\hat{m}^2)} \end{aligned} \quad (105)$$

- **Polynomial invariants**

Superpolynomial:

$$P_{4_1}(a, q, t) = P_0(a, q, t) [a^{-1} t^{-2} + q^{-1} t^{-1} + 1 + qt + at^2] \quad (106)$$

B4. Knot 5_2

- **Representation through quantum R -matrix and Drinfeld associator**

The polynomial invariant for 5_2 can be constructed using formulae (15) and (18). In this case, the knot can be represented as the closure of the following element:

$$b_{5_2} = g_2^3 g_1 g_2^{-1} g_1 \in B_3 \quad (107)$$

In this representation, the knot has 5 positively oriented crossings and one negatively oriented ($w(b) = 4$) and two strings ($n = 3$ in (18)).

- **Representation through Vassiliev invariant**

The first Vassiliev invariants of knot 5_2 are:

$\alpha_{2,1}$	$\alpha_{3,1}$	$\alpha_{4,1}$	$\alpha_{4,2}$	$\alpha_{5,1}$	$\alpha_{5,2}$	$\alpha_{5,3}$
8	-24	$\frac{268}{3}$	$\frac{44}{3}$	-368	-64	-56

(108)

For instance, the usage of (21) for $SU(2)$ gives:

$$\langle 5_2 \rangle = [N]_q \exp \left(-8J(1+J)h^2 + 24J(1+J)h^3 + \frac{4}{3}J(1+J)(-67 + 22J + 22J^2)h^4 + \dots \right) \quad (109)$$

with $h = 2\pi i \hbar$.

- **Representation through quantum dilogarithm**

The colored Jones polynomial for 5_2 with $N = 2J + 1$ can be represented in the hypergeometric form:

$$K_N(5_2|q) = [N]_q \sum_{0 \leq k \leq l \leq N-1} \frac{(q, N)_l (q, N)_l}{(q, N)_k^*} q^{-k(l+1)} \equiv [N]_q J_N(5_2|q) \quad (110)$$

Then, (63) gives:

$$J_N \sim \sum_{0 \leq k \leq l \leq N-1} (-i)^{k+2l} q^{-k(l+1)} \frac{s \left(i\epsilon_2(N-1-l) + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)^2 s \left(i\epsilon_2(N+k) + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)}{s \left(i\epsilon_2(N-1) + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)^2 s \left(i\epsilon_2 N + \frac{i}{2}(\epsilon_1 + \epsilon_2) \middle| \epsilon_1, \epsilon_2 \right)}$$

$$J_N = \frac{i}{s(u)^3} \int dz_1 \int dz_2 s(u-z \middle| \epsilon_1, \epsilon_2)^2 s(u+z \middle| \epsilon_1, \epsilon_2) e^{-\frac{2\pi z_1 z_2}{\epsilon_1 \epsilon_2} + \frac{\pi i(z_1 - z_2)}{\epsilon_1} + \frac{2\pi i(2z_1 + z_2)}{\epsilon_2}} \quad (111)$$

Again, the presence of three dilogarithms in the integral (111) implies that the space $S^3 \setminus 5_2$ can be realized by gluing three tetrahedrons in the Hikami state model.

- **$\mathcal{A}$ -polynomial**

In this case, one has

$$\mathcal{A}(l, m) = 1 + l(-1 + 2m + 2m^2 - m^4 + m^5) + l^2(m^2 - m^3 + 2m^5 + 2m^6 - m^7) + l^3 m^7 \quad (112)$$

The quantum $\mathcal{A}$ -polynomial can be calculated as in the previous cases, the result reads

$$\begin{aligned} \hat{\mathcal{A}}(l, m) = & q^{\frac{1}{2}}(1 - q^4 \hat{m}^2)(1 - q^5 \hat{m}^2) - (1 - q^2 \hat{m}^2)(1 - q^5 \hat{m}^2)(1 - 2q\hat{m} - q(q + q^3)\hat{m}^2 + \\ & + q^2(1 - q)(1 - q^2)\hat{m}^3 + q^5 \hat{m}^4 - q^6 \hat{m}^5) \hat{l} + q^{\frac{5}{2}}(1 - q\hat{m}^2)(1 - q^4 \hat{m}^2)\hat{m}^2(1 - q^2 \hat{m} - q^2(1 - q)(1 - q^2)\hat{m}^2 + \\ & + q^4(1 + q^3)\hat{m}^3 + 2q^7 \hat{m}^4 - q^9 \hat{m}^5) \hat{l}^2 + q^{14}(1 - q\hat{m}^2)(1 - q^2 \hat{m}^2)\hat{m}^7 \hat{l}^3 \end{aligned}$$

This is the simplest example, when $\hat{\mathcal{A}}(l, m)$ is not quadratic, but cubic in l . In this case the spectral curve is not hyperelliptic.

- **Polynomial invariants**

Superpolynomial:

$$P_{5_2}(a, q, t) = P_0(a, q, t) [aq^{-1} + at + aqt^2 + a^2 q^{-1} t^2 + a^2 t^3 a^2 q t^4 + a^3 t^5] \quad (113)$$

Appendix C. Examples of the volume conjecture

The volume conjecture states that, at large N , logarithm of the colored Jones polynomial of the hyperbolic knot K is proportional to the volume of the knot complement $S^3 \setminus K$:

$$\log |J_N(K)| \sim \frac{N}{2\pi} \text{Vol}(S^3 \setminus K) \quad \text{for } q = e^{2\pi i/N}, \text{ and } J_N(K) = \langle K \rangle / \dim_q(R) \quad (114)$$

Here we represent the plots of $\log |J_N(K)|$ for $N = 1..100$, for the hyperbolic knot 4_1 and for the toric knot 3_1 .

For the knot 4_1 , formula (100) for $q = e^{2\pi i/N}$ gives

$$J_N(4_1) = \sum_{k=1}^{N-1} \prod_{j=1}^k 4 \sin^2 \frac{\pi j}{N} \quad (115)$$

Note that all the terms in this formula are positive. Using representation (99) and the expansion of quantum dilogarithm function (45) at large N , this sum might be approximated by the integral

$$J_N(4_1) \sim \int dz \exp \left(\frac{N}{2\pi i} (\text{Li}_2(-e^{-iz}) - \text{Li}_2(-e^{iz})) \right) \quad (116)$$




Figure 3: The plot of $\ln J_N(4_1)$ for $N = 3 \dots 100$. At large N the plot becomes linear.

Using the saddle point approximation, one obtains

$$\frac{d}{dz}(\text{Li}_2(-e^{-iz}) - \text{Li}_2(-e^{iz})) = 0 \Rightarrow \log((1 + e^{-iz})(1 + e^{iz})) = 0 \Rightarrow e^{iz} = e^{\frac{2\pi i}{3}} \quad (117)$$

and for the volume one gets

$$\text{Vol}(4_1) = -i \left(\text{Li}_2(-e^{-\frac{2\pi i}{3}}) - \text{Li}_2(-e^{\frac{2\pi i}{3}}) \right) \approx 2.02688 \quad (118)$$

One can directly measure this value as a slope in the plot, Fig.3.

In the case of the toric knot 3_1 , the quantity $J_N(3_1)$ may take complex values, and, therefore, the saddle point approximation is more subtle. Fig.4 shows the power behaviour of the absolute value asymptotics which is expected [55] and is consistent with the volume conjecture for the torus knot

$$|J_N(3_1)| \sim N^{3/2}$$

while the phase of $J_N(3_1)$ behaves in a much more tricky way [56].




Figure 4: The plot of $\ln |J_N(3_1)|$ for $N = 3 \dots 100$ as a function of $\ln N$. At large N the behavior of the plot is linear with the slope $3/2$.

References

- [1] Yu.Terashima and M.Yamazaki, arXiv:1103.5748
- [2] D.Gaiotto, arXiv:0904.2715
- [3] D. Gaiotto, G. Moore, A. Neitzke, arXiv:0907.3987;
T. Dimofte, S. Gukov, Y. Soibelman, Lett.Math.Phys. **95** (2011) 1-25, arXiv:0912.1346;
M. Kontsevich and Y. Soibelman, 0811.2435;
V. Fock and A. Goncharov, Publ. Math. Inst. Hautes Études Sci. **103** (2006) 1-211, math/0311149
- [4] V.Pestun, arXiv:0906.0638
- [5] N.Nekrasov and E.Witten, JHEP 1009 (2010) 092, arXiv:1002.0888
- [6] H.Awata and H.Kanno, arXiv:0910.0083
- [7] E.Gorsky, arXiv:1003.0916
- [8] T.Dimofte, S.Gukov and L.Hollands, arXiv:1006.0977
- [9] R.Dijkgraaf, H.Fuji and M.Manabe, arXiv:1010.4542
- [10] K. Hosomichi, S. Lee and J. Park, JHEP **1012** (2010) 079, arXiv:1009.0340; JHEP **1103** (2011) 127, arXiv:1012.3512; arXiv:1102.4716
- [11] E.Witten, arXiv:1001.2933; arXiv:1101.3216
- [12] F.Dolan, V.Spiridonov and G.Vartanov, arXiv:1104.1787
- [13] L.Alday, D.Gaiotto and Y.Tachikawa, Lett.Math.Phys. **91** (2010) 167-197, arXiv:0906.3219;
N.Wyllard, JHEP **0911** (2009) 002, arXiv:0907.2189;
A.Mironov and A.Morozov, Phys.Lett. **B680** (2009) 188-194, arXiv:0908.2190; Nucl.Phys. **B825** (2009) 1-37, arXiv:0908.2569

- [14] K.Hikami, Int.J.Mod.Phys. **A16** (2001) 3309-3333, math-ph/0105039; Int.J.Mod.Phys. **B16** (2002) 1963-1970; J.Geom.Phys. **57** (2007) 1895-1940, math/0604094;
T. Dimofte, S. Gukov, J. Lenells, D. Zagier, Commun.Num.Theor.Phys. **3** (2009) 363-443, arXiv:0903.2472
- [15] R.Poghossian, JHEP **0912** (2009) 038, arXiv:0909.3412;
V.Alba and And.Morozov, JETP Lett. **90** (2009) 708-712, arXiv:0911.0363
- [16] J. Teschner, Int.J.Mod.Phys. **A19S2** (2004) 436-458, hep-th/0303150; Contribution to 14th International Congress on Mathematical Physics (ICMP 2003), hep-th/0308031
- [17] E.W.Barnes, Proc. London Math. Soc. **31** (1899) 358-381; Phil. Trans. Roy. Soc. **A196** (1901) 265-387; Trans. Cambr. Phil. Soc. **19** (1904) 374-425
- [18] T.Shintani, J. Fac. Sci. Univ. Tokyo, Sect. 1A **24** (1977) 167-199;
N.Kurokawa, Proc. Japan Acad. **A67** (1991) 61-64; Proc. Japan Acad. **A68** (1992) 256-260; Adv. Studies Pure Math. **21** (1992) 219-226;
L.Faddeev and R.Kashaev, Mod.Phys.Lett. **A9** (1994) 427-434, hep-th/9310070;
L.Faddeev, Lett. Math. Phys. **34** (1995) 249; hep-th/9504111; math.qa/9912078;
S.Sergeev, V.Bazhanov, H.Boos, V.Mangazeev and Yu.Stroganov, IHEP-95-129, 1995;
M.Jimbo and T.Miwa, J.Phys. **A29** (1996) 2923-2958; hep-th/9601135;
S. Ruijsenaars, J. Math. Phys. **38** (1997) 1069-1146;
R. Kashaev, Lett.Math.Phys. **43** (1998) 105-115
- [19] E.Witten, Comm.Math.Phys. **121** (1989) 351-399
- [20] J.Wess and B.Zumino, Phys.Lett. **B37** (1971) 95;
S.Novikov, UMN, **37** (1982) 37;
E.Witten, Comm.Math.Phys. **92** (1984) 455;
A.Gerasimov, A.Marshakov, A.Morozov, M.Olshanetsky, S. Shatashvili, Int.J.Mod.Phys. **A5** (1990) 2495-2589
- [21] A.Morozov, Phys.Usp.(UFN) **35** (1992) 671-714; **37** (1994) 1, hep-th/9303139; hep-th/9502091; hep-th/0502010;
A.Mironov, Int.J.Mod.Phys. **A9** (1994) 4355, hep-th/9312212; Phys.Part.Nucl. **33** (2002) 537; hep-th/9409190; hep-th/9409190; Theor.Math.Phys. **114** (1998) 127, q-alg/9711006
- [22] N.M.Dunfield, S.Gukov and J.Rasmussen, Experimental Math. **15** (2006) 129-159, arXiv:math/0505662
- [23] S.Gukov, A.Iqbal, C.Kozcaz and C.Vafa, arXiv:0705.1368
- [24] M. Khovanov, Duke Math. J. **101** (2000) 359-426
- [25] M. Aganagic, R. Dijkgraaf, A. Klemm, M. Marino, C. Vafa, Commun.Math.Phys. **261** (2006) 451-516, hep-th/0312085;
M.Marino, Rev.Mod.Phys. **77** (2005) 675-720, hep-th/0405083
- [26] S.Garoufalidis and T.Le, Geometry and Topology, **9** (2005) 1253-1293, math/0309214
- [27] A.Alexandrov, A.Mironov and A.Morozov, Int.J.Mod.Phys. **A19** (2004) 4127, hep-th/0310113; Teor.Mat.Fiz. **150** (2007) 179-192, hep-th/0605171; Physica **D235** (2007) 126-167, hep-th/0608228; JHEP **12** (2009) 053, arXiv:0906.3305;
A.Alexandrov, A.Mironov, A.Morozov, P.Putrov, Int.J.Mod.Phys. **A24** (2009) 4939-4998, arXiv:0811.2825;
B.Eynard, JHEP **0411** (2004) 031, hep-th/0407261;
L.Chekhov and B.Eynard, JHEP **0603** (2006) 014, hep-th/0504116; JHEP **0612** (2006) 026, math-ph/0604014;
N.Orantin, arXiv:0808.0635
- [28] E. Guadagnini, M. Martellini, M. Mintchev, In "Clausthal 1989, Proceedings, Quantum groups", 307-317; Phys.Lett. **B235** (1990) 275
- [29] A.Morozov and A.Smirnov, Nucl.Phys. **B835** (2010) 284-313, arXiv:1001.2003
A.Smirnov, hep-th/0910.5011, to appear in the Proceedings of International School of Subnuclear Phys. in Erice, Italy, 2009

- [30] V.Drinfeld, Published in "Problems of modern quantum field theory" 1-13, Alushta, 1989
- [31] T.T.Q.Le and J. Murakami, Nagoya Math. J.Vol. **142** (1996) 39-65
- [32] S.Chmutov and S.Duzhin, math/0501040
- [33] V.Knizhnik and A.Zamolodchikov, Nucl.Phys. **B247** (1984) 83-103
- [34] P. Dunin-Barkovsky, A. Sleptsov and A. Smirnov, to appear
- [35] S. Chmutov, S. Duzhin, J. Mostovoy, arXiv:1103.5628;
J.M.F. Labastida, E. Perez, J.Math.Phys. **41** (2000) 2658-2699, hep-th/9807155 J.Math.Phys. **39** (1998) 5183-5198, hep-th/9710176;
M. Alvarez, J.M.F. Labastida, E. Perez, Nucl.Phys. **B488** (1997) 677-718, hep-th/9607030;
M. Alvarez, J.M.F. Labastida, Commun.Math.Phys. **189** (1997) 641-654, q-alg/9604010; Nucl.Phys. **B433** (1995) 555-596, hep-th/9407076
- [36] Dror Bar-Natan, Edward Witten, Commun.Math.Phys. **141** (1991) 423-440;
Dror Bar-Natan, Ph.D. Thesis; J.Knot Theor.Ramifications **4** (1995) 503-547; q-alg/9703025;
R. Kashaev and N.Reshetikhin, math/0202211; math/0202212;
V.G. Turaev, O.Y. Viro, Topology **31** (1992) 865-902;
V. G. Turaev, UMN, **41** (247) (1986) 97-147;
S. Axelrod, I. Singer, Proc. XXth Internat. Conf. on Differential Geometric Methods in Theoretical Physics, arXiv:hep-th/9110056v1
- [37] M. Goussarov, M. Polyak, O. Viro, Topology, **39(5)** (2000) 1045-1068
- [38] A.Mironov, A.Morozov and S.Natanzon, Theor.Math.Phys. **166** (2011) 1-22, arXiv:0904.4227; arXiv:1012.0433
- [39] R.Kashaev, arXiv:q-alg/9601025
- [40] M.Marino, JHEP **0812** (2008) 114, arXiv:0805.3033
- [41] A.Morozov and Sh.Shakirov, JHEP **0912** (2009) 003 arXiv:0906.003; arXiv:1007.4100
- [42] W.Turston, *The Geometry and Topology of Three-Manifolds*, Lecture Notes at Princeton University, 1980
- [43] R.Gelca, Math. Proc. Cambridge Philos. Soc. **133** (2002) 311-323;
R. Gelca and J. Sain, J. Knot Theory Ramifications **12** (2003) 187-201
- [44] S. Garoufalidis and Thao Vuong, arXiv:1010.3147;
S. Garoufalidis and C. Koutschan, arXiv:1011.6329
- [45] N.Nekrasov, Nucl.Phys. **B531** (1998) 323-344, arXiv:hep-th/9609219;
A.Gorsky, S.Gukov and A.Mironov, Nucl.Phys., **B518** (1998) 689, arXiv:hep-th/9710239;
A.Marshakov, A.Mironov, Nucl.Phys., **B518** (1998) 59-91, arXiv:hep-th/9711156;
H.W. Braden, A.Marshakov, A.Mironov and A.Morozov, Phys.Lett., **B448** (1999) 195, hep-th/9812078;
Nucl.Phys., **B558** (1999) 371, hep-th/9902205
- [46] A.Mironov and A.Morozov, JHEP **04** (2010) 040, arXiv:0910.5670; J.Phys. **A43** (2010) 195401, arXiv:0911.2396
- [47] A.Popolitov, arXiv:1001.1407;
Wei He, Yan-Gang Miao, Phys.Rev. **D82** (2010) 025020, arXiv:1006.1214;
K.Maruyoshi and M.Taki, arXiv:1006.4505;
A.Marshakov, A.Mironov and A.Morozov, arXiv:1011.4491;
F.Fucito, J.F.Morales, R.Poghossian and D. Ricci Pacifici, arXiv:1103.4495;
Y.Zenkevich, arXiv:1103.4843;
N.Dorey, T.J.Hollowood and S.Lee, arXiv:1103.5726
- [48] N.Nekrasov and S.Shatashvili, arXiv:0908.4052
- [49] A.Mironov, A.Morozov, A.Smirnov and Sh.Shakirov, *Proving AGT conjecture for $\beta = 1$: extension to five dimensions*, to appear

- [50] S.Kharchev, D.Lebedev and M.Semenov-Tian-Shansky, Commun.Math.Phys. **225** (2002) 573-609, arXiv:hep-th/0007040
- [51] N.Seiberg and E.Witten, Nucl.Phys., **B426** (1994) 19-52, hep-th/9408099; Nucl.Phys., **B431** (1994) 484-550, hep-th/9407087;
A.Gorsky, I.Krichever, A.Marshakov, A.Mironov, A.Morozov, Phys.Lett., **B355** (1995) 466-477, hep-th/9505035;
R.Donagi and E.Witten, Nucl.Phys., **B460** (1996) 299-334, hep-th/9510101
- [52] H.Ooguri and C.Vafa, Nucl.Phys. **B577** (2000) 419, hep-th/9912123
- [53] T. Dimofte, arXiv:1102.4847
- [54] R. Dijkgraaf and H. Fuji, Fortsch.Phys. **57** (2009) 825-856, arXiv:0903.2084
- [55] R.Kashaev and O.Tirkkonen, arXiv:math/9912210
- [56] H.Murakami, Internat. J. Math. **15** (2004) 547-555, arXiv:math/0405126